


Southern Pine Beetle Historical Data for Georgia Georgia's Most Destructive Forest Pest

	Year	Total Spots	\$ loss
	1972	3,766	\$970,771
	1973	4,263	\$6,290,700
	1974	12,752	\$7,081,540
	1975	7,201	\$1,324,985
	1976	2,300	\$620,625
36 year totals	1977	1,388	\$306,777
	1978	1,221	\$172,468
# Spots	1979	22,254	\$25,216,776
	1980	19,765	\$19,737,886
\$ Loss	1981	507	\$95,092
	1982	955	\$849,789
	1983	1,906	\$1,763,285
	1984	61	\$438,360
	1985	1,278	\$4,879,991
	1986	8,426	\$9,673,712
	1987	2,215	\$2,302,346
	1988	6,485	\$14,111,039
	1989	2,043	\$8,405,832
36 year averages:	1990	157	\$3,005,106
	1991	3,606	\$3,787,367
# Spots	1992	5,210	\$3,515,585
	1993	1,583	\$2,916,690
\$ Loss	1994	1,084	\$771,098
	1995	5,750	\$27,866,516
	1996	279	\$5,681,934
	1997	1,023	\$2,609,742
	1998	549	\$1,943,721
	1999	147	\$1,472,332
	2000	1,798	\$8,578,988
	2001	2,438	\$14,292,363
	2002	9,070	\$57,239,375
	2003	239	\$931,899
	2004	79	\$370,116
	2005	0	\$0
	2006	601	\$928,545
	2007	1,704	\$14,141,750



For more information on the southern pine beetle, visit: <http://whizlab.isis.vt.edu/servlet/sf/spbcc/index.html>

For more information about the data contained in this report, contact James Johnson (706) 542-9608 jjohnson@gfc.state.ga.us

Based upon annual aerial surveys conducted by the Georgia Forestry Commission