



2024 Georgia Southern Pine Beetle Prediction Survey Report

Georgia Forestry Commission
Forest Health Staff

The Georgia Forestry Commission (GFC) participates annually in the southern pine beetle (SPB) prediction trapping program. The southern pine beetle is the most destructive forest pest in the southeastern states. Survey results are documented in an annual report so that activity and damage levels can be anticipated and mitigated.

The Georgia Forestry Commission follows the SPB Prediction Trapping protocol set up by Texas A&M. A 12-funnel Lindgren trap is baited with a three lure system: Frontalin, Sirex and the endo-brevicomin, flexlure (Figure 1). Traps are placed in the field to coincide with redbud bloom (around the first of March in the southern part of the state and mid-March in the northern part of the state). The Georgia Forestry Commission placed 50 traps across the state in 50 counties and six weekly samples were collected from each trap (Figure 2). The number of SPB and clerid beetles (a natural predator of SPB) were counted each week. For 2024, the average number of SPB per-trap per-day was **41.0** (Figure 3). This number is considered in the moderate range. Nine counties, Carroll, Clarke, Dawson, Greene, Heard, Morgan, Oconee, Putnam and Stephens, had SPB per-trap per-day counts greater than 81. These counties are in the high range for SPB per-trap per day. Six counties, Jasper, Macon, Meriwether, Monroe, Oglethorpe and Twiggs, had SPB per-trap per-day counts greater than 50 but fewer than 80. Ten counties, Baldwin, Floyd, Harris, Jones, Laurens, Polk, Rabun, Randolph, Stewart and Webster had SPB per-trap per-day counts greater than 20 but fewer than 50. All these counties are in the moderate range for SPB per-trap per-day. Dawson County had the highest overall SPB per-trap per-day catch at **203.2**, placing it in the high category. The remaining 26 traps had an average of fewer than 20 SPB per-trap per-day.



Figure 1: Lindgren funnel trap used to monitor southern pine beetle populations.

In 2018, the USDA Forest Service (USFS) and all southeastern state cooperators collaborated with Dartmouth College and Bates College, through the Science and Technology Development Program, (STDP) to develop a new prediction model. They found the two greatest factors in predicting the probability of the area having SPB spots are the number of SPB collected per two-week period in the current year and the number of SPB spots in that county the previous year. This year is the sixth year GFC is using this model. The results from the 2024 survey predict that overall SPB activity will be low in the southern region and moderate in the western and north regions of Georgia. The central region has five areas having a high probability (>50%) of any spots. Greene, Jasper, Jones and Oglethorpe Counties have high probability of having any spots with Putnam County having the greatest probability.

Perhaps a more informative prediction, because it accounts for more variability within the historical data, is the probability of greater than 50 spots in a county (Figures 4&5). Putnam County has the highest probability of having greater than 50 spots, with a 43.7% chance. Greene, Jasper, Jones and Oglethorpe Counties have a 25.0-37.3% chance of having greater than 50 spots. Clarke, Dawson, Macon, Morgan, Oconee, Stephens and Wilkinson have a 5-12.9% chance of having greater than 50 spots. The remaining 38 counties that were trapped in 2024 have 4.5% or less chance of having greater than 50 spots. To put this in perspective, over the last 13 years, we have only seen the total number of spots statewide rise above 50 four out of 13 years.

These prediction models help guide landowners in management decisions. They do not guarantee that outbreaks will or will not occur on their property. The best advice is for landowners to manage for healthy forests with techniques such as thinning, prescribed burning, and invasive species control. In the past decade, SPB outbreaks have been limited to infestations in stands that are either over-stocked or over-mature. Southern pine beetle favors pine stands that are over-crowded and stressed. The Georgia Forestry Commission will continue to monitor locations of beetle spots throughout the year. All reported beetle activity will be surveyed and monitored to mitigate damage for landowners.

Routine annual aerial surveys in summer 2024 will be conducted to document and monitor pine beetle activity. Scheduled flights will be conducted across the state following predetermined flight lines. Aerial surveys can also be conducted by the GFC Air Operations Division during normal flights and any new possible infestations found from these flights will be investigated using ground surveys. All infestations will be reported to landowners, and GFC foresters will work with landowners to limit damage and control infestations.

For additional information contact:

Michael Torbett
Forest Health Coordinator
Georgia Forestry Commission
mtorbett@gfc.state.ga.us

Mark McClure
Forest Health Specialist
Georgia Forestry Commission
mmcclure@gfc.state.ga.us




Figure 2: SPB trapping counties 2024.




Figure 3: Southern Pine Beetle trap locations and the number of SPB caught per day over the six-week period.




Figure 4: Probability of 50 or more spots




Figure 5: Probability of 50 or more spots on a regional scale